

## ROLE OF HRCT THORAX IN TRACHEOBRONCHIAL FOREIGN BODY DETECTION

Manphool Singh Maharia<sup>1</sup>, Sandeep Kumar<sup>2</sup>, Deep Chand<sup>3</sup>, Gaurav Gupta<sup>4</sup>

<sup>1</sup>Senior Resident, <sup>2</sup>Resident, <sup>3</sup>Senior Professor and Head, <sup>4</sup>Associate Professor

<sup>1,3,4</sup> Department of ENT, Sardar Patel Medical College, Bikaner, Rajasthan, India

<sup>2</sup> Department of Radiology, RNT Medical College, Udaipur, Rajasthan, India

**Article Info:** Received 10 February 2019; Accepted 11 March. 2019

**Cite this article as:** Maharia, M., Kumar, S., Chand, D., & Gupta, G. (2019). ROLE OF HRCT THORAX IN TRACHEOBRONCHIAL FOREIGN BODY DETECTION. *International Journal of Medical and Biomedical Studies*, 3(3).

**DOI:** <https://doi.org/10.32553/ijmbs.v3i3.144>

**Address for Correspondence:** Sandeep Kumar, Department of Radiology, RNT Medical College, Udaipur, Rajasthan, India

**Conflict of interest:** No conflict of interest.

### Abstract

**Background:** Aspirated foreign bodies in the airway continue to present challenges to the otorhinolaryngologist and interventional pulmonologists. To established the role of HRCT in diagnosis of foreign bodies in suspected cases.

**Methods-**30 patients with suspected FB inhalation on the basis of clinical history and symptoms like respiratory distress, stridor, and history of choking were recruited for study in department of Otorhinolaryngology in Sardar Patel Medical College and Hospital, Bikaner Rajasthan.

**Results:** 29 patients (96.67%) with foreign bodies were identified on chest CT. For the patients with tracheobronchial foreign bodies, the occurrence of unilateral hyperlucent lung and post-obstructive lobar or segmental infiltrates on plain chest X-ray was 43.33%. 17 patients (56.67%) had no abnormalities on plain X-ray. The difference between multidetector CT and plain X-ray results was statistically significant.

**Conclusion:** Foreign bodies are missed by clinical and X-ray examination in many cases and that is only picked up by HRCT. Thus it is the ideal modality in diagnosis of foreign bodies to avoid the morbidity and mortality associated with missing foreign body.

**Keywords:** foreign bodies, high resolution computed tomography, X-ray

### Introduction:

Aspirated foreign bodies in the airway continue to present challenges to the otorhinolaryngologist and interventional pulmonologists. Inhalation of foreign bodies is common in pediatric age group and 94 % of them occur in infants and children, with peak incidence in the age group of 1-3 years and is very rare in adults. <sup>1</sup> Adults frequently have an underlying condition associated with impairment of airway

protection such as mental retardation, neurological disorder, alcohol or sedative abuse.  
<sup>2-4</sup>

The symptoms and signs produced depend upon the size, nature, location and area of lodgment of the foreign body in the trachobronchial tree. A large foreign body occluding the upper airway may lead to sudden death, whereas a small foreign body lodged in the bronchial tree may present with less severe system.

The radiological diagnosis of FB inhalation is challenging for several reasons. Chest radiography may show a variety of findings including airtrapping, consolidation, and atelectasis and bilateral over aeration. Only 10 % of FB is radiopaque. The findings of chest radiography are normal in up to 30% of children who inhaled a FB and the presence of pulmonary infiltrates may misdirect the management of FB inhalation.<sup>5</sup> Bronchoscopy is often performed for definitive diagnosis and management, however, it is invasive and procedure related serious complications may occur. Recently developed high resolution computed tomography (HRCT) and virtual bronchoscopy is a noninvasive technique that provides realistic 3D views of the tracheobronchial tree. In addition to the detection of foreign body HRCT and virtual bronchoscopy can help the surgeon plan for operative bronchoscopy and safe removal of foreign body.<sup>6-7</sup>

**Material and method**

30 patients with suspected FB inhalation on the basis of clinical history and symptoms like respiratory distress, stridor, and history of choking were recruited for study in department of Otorhinolaryngology in Sardar Patel Medical College and Hospital, Bikaner Rajasthan. These patients were further underwent chest X-Ray and findings were noted followed by HRCT was performed in all patients and images were taken. The presence of FB, its location, size and density were determined by consultant radiologist. Associated findings i.e. collapse, consolidation; emphysema, mediastinal shift and pneumothorax were also noted. The finding in form of age, sex, type and site of FB noted and results of HRCT compare with X-ray and bronchoscopy.

**Results**

**Table 1: Socio-demographic variable**

Socio-demographic variable		No of cases	Percentage
Age	< 1 yrs	3	10.00
	1-3 yrs	16	53.33
	3-6 yrs	9	30.00
	>6 yrs	2	6.66
Sex	Male	19	63.33
	Female	11	36.67

Maximum patients (53.33%) were 1-3 yrs age group and 63.33% patients were male.

**Table 2: Diagnosis wise distribution**

Diagnosis		No. of cases	Percentage
X-ray chest	Identified FB	13	43.33%
	Not Identified FB	17	56.67%
HRCT chest	Identified FB	29	96.67%
	Not Identified FB	1	3.33%

29 patients (96.67%) with foreign bodies were identified on chest CT. For the patients with tracheobronchial foreign bodies, the occurrence of unilateral hyperlucent lung and post-obstructive lobar or segmental infiltrates on plain chest X-ray was 43.33%. 17 patients (56.67%) had no abnormalities on plain X-ray. The difference between multidetector CT and plain X-ray results was statistically significant.

### Discussion

The foreign body into the tracheobronchial tree occurs in all age groups, infants and small children suffer most commonly. The anatomic relation of the larynx, shouting, crying and playing while eating and lack of parental supervision contributes to this hazard. Most patients in the present study were below 3 years and smallest child was 6 month old with male to female ratio was 2:1 which is similar to the that reported in other study.<sup>8</sup>

In our study for the patients with tracheobronchial foreign bodies, the occurrence of unilateral hyperlucent lung and post-obstructive lobar or segmental infiltrates on plain chest X-ray was 43.33%. 17 patients (56.67%) had no abnormalities on plain X-ray. and these findings supported by other study<sup>9</sup> Sensitivity of chest X-ray in foreign body detection is low in our study although it is much lesser than the previous studies thus, although chest radiography may help, it seems neither sufficiently sensitive nor specific for the diagnosis of foreign body aspiration.<sup>10,11</sup> 29 patients (96.67%) with foreign bodies were identified on chest CT. Multidetector CT scan chest is the diagnostic technique used for detection of foreign bodies. It not only can reveal foreign bodies in the bronchial tree but also is very sensitive in detecting associated findings<sup>12</sup>

### Conclusions

The diagnosis of foreign body aspiration of the airway in children can be accomplished by using chest HRCT. It is often useful in delineating the exact shape, location, volume and form of a bronchial foreign body and can help the surgeon

plan for operative bronchoscopy and safe removal of the foreign body.

### Bibliography

1. Rothman BF, Boeckman CR. Foreign bodies in the larynx and tracheobronchial tree in children. A review of 225 cases, *Annal Otol, Rhinol laryngol.* 1980;89:434-6.
2. Prakash UBS, Cortese DA. Tracheobronchial foreign bodies. In Prakash UBS, ed. *Bronchoscopy.* New York Raven Press. 1994:253-78.
3. Limper AH, Prakash UBS. Tracheobronchial bodies in adults. *Ann Intern Medicine.* 1990;112:604-9.
4. Chen CH, Lai CL, Tsai TT, Lee YC, Perng RP. Foreign body aspiration into lower airway in Chinese adults. *Chest.* 1997;112:129-33.
5. Khan MF, Herzog C, Ackermann H et al (2004) Virtual endoscopy of the tracheobronchial system: sub-millimeter collimation with the 16- row multidetectorscanner. *EurRadiol* 14:1400– 5
6. Tan HKK, Brown K, McGill T, Kenna MA, Lund DP, Healy GB. Airway foreign bodies (FB): a 10- year review. *Int J Pediatr Otorhinolaryngol* 2000; 56:91 –9
7. Finkelstein SE, Schrupp DS, Nguyen DM et al (2003) Comparative evaluation of super higher solution CT scan and virtual bronchoscopy for the detection of tracheobronchial malignancies. *Chest* 124:1834–40
8. Kalyanappagol VT, Kulkarni NH, Bidri LH. Management of tracheobronchial foreign body.aspirations in paediatric age group - a 10 year retrospective analysis. *Indian J Anaesth.* 2007;51:23– 27.
9. Rogalla P, Stöver B, Scheer I, Juran R, Gaedicke G, Hamm B. Low- dose spiral CT: applicability to pediatric chest imaging. *Pediatr Radiol* 1998;28 : 365–69

10. Zwirewich CV, Mayo JR, Muller NL. Low-dose high-resolution CT of lung parenchyma. *Radiology* 1991; 180:413 –17
11. Black RE, Johnson DG, Matlak ME. Bronchoscopic removal of aspirated foreign bodies in children. *J PadiatrSurg* 1994; 29:682-4.
12. James B, Snow JR. Bronchology. In *Disease of the nose, throat, ear, head and neck*, volume 2, 14th Ed. 1991; 1278-96