

Study of Serum Adiponectin and Leptin Levels in Patients with Type 2 Diabetes Mellitus

Srushtee Bipin Jibhkate

Associate Professor, Department of Biochemistry, Naraina Medical College & Research Centre,
Panki, Kanpur Uttar Pradesh

Article Info: Received 16 August 2022; Accepted 19 September 2022

Corresponding author: Srushtee Bipin Jibhkate

Conflict of interest: No conflict of interest.

Abstract

Type 2 diabetes mellitus (T2DM) is a global health problem characterized by insulin resistance and metabolic dysfunction. Adiponectin and leptin, two key adipokines, play a crucial role in energy homeostasis, glucose metabolism, and insulin sensitivity. This study aimed to evaluate serum adiponectin and leptin levels in T2DM patients and compare them with healthy controls.

A case-control study was conducted with 100 participants, including 50 T2DM patients and 50 healthy controls. Serum adiponectin and leptin levels were measured using ELISA kits, and their correlations with metabolic parameters such as fasting blood glucose (FBG), HbA1c, and body mass index (BMI) were analyzed. The results showed significantly lower adiponectin and higher leptin levels in T2DM patients compared to controls ($p < 0.05$). Additionally, adiponectin showed a negative correlation with FBG and HbA1c, while leptin levels correlated positively with BMI and insulin resistance.

The findings suggest that adiponectin deficiency and leptin excess contribute to insulin resistance and metabolic dysregulation in T2DM patients. These adipokines could serve as potential biomarkers for assessing diabetes risk and complications. Further research is required to explore therapeutic interventions targeting adipokine modulation for improved glycemic control.

Keywords: Type 2 diabetes mellitus, adiponectin, leptin, insulin resistance, metabolic syndrome, adipokines.

Introduction

Type 2 diabetes mellitus (T2DM) is a chronic metabolic disorder characterized by insulin resistance, hyperglycemia, and progressive β -cell dysfunction. The global prevalence of T2DM has increased significantly, mainly due to sedentary lifestyles, obesity, and genetic predisposition [1]. The dysregulation of adipokines, particularly adiponectin and leptin, plays a crucial role in the pathophysiology of diabetes and its associated complications [2].

Adiponectin is an insulin-sensitizing adipokine secreted primarily by adipocytes. It has anti-inflammatory, anti-atherogenic, and insulin-sensitizing properties, making it a key regulator of glucose metabolism [3]. Low adiponectin levels are commonly observed in T2DM and obesity, which contribute to insulin resistance

and cardiovascular risk [4]. Adiponectin enhances glucose uptake by muscles, suppresses hepatic gluconeogenesis, and promotes fatty acid oxidation [5]. Several studies indicate a negative correlation between adiponectin and insulin resistance, making it a potential therapeutic target for diabetes management [6].

Leptin, another key adipokine, is responsible for regulating energy balance and appetite control. It is secreted in proportion to body fat mass and acts on the hypothalamus to suppress hunger [7]. However, in obesity and T2DM, leptin resistance develops, leading to hyperleptinemia and persistent metabolic dysregulation [8]. High leptin levels are associated with inflammation, endothelial dysfunction, and insulin resistance,

making it a potential biomarker for T2DM progression [9].

The adiponectin-leptin ratio is considered an indicator of insulin sensitivity, with lower ratios predicting higher metabolic risk [10]. Understanding the role of adipokines in T2DM could provide insights into potential therapeutic strategies aimed at modulating their levels. This study aims to evaluate serum adiponectin and leptin levels in T2DM patients and analyze their correlations with glucose metabolism and obesity-related markers.

Aim and Objectives

Aim

To evaluate serum adiponectin and leptin levels in patients with Type 2 diabetes mellitus and compare them with healthy controls.

Objectives

1. To assess serum adiponectin and leptin levels in T2DM patients and healthy individuals.
2. To analyze the correlation between adiponectin, leptin, and metabolic parameters such as fasting blood glucose, HbA1c, and BMI.

Table 1: Serum Adiponectin and Leptin Levels in T2DM Patients and Controls

Parameter	T2DM Patients (Mean ± SD)	Controls (Mean ± SD)	p-Value
Adiponectin (µg/mL)	3.4 ± 1.2	7.8 ± 2.1	<0.001
Leptin (ng/mL)	19.6 ± 5.3	9.2 ± 3.1	<0.001

Table 2: Correlation Between Adipokines and Metabolic Parameters

Parameter	Adiponectin (r-value)	Leptin (r-value)	p-Value
FBG (mg/dL)	-0.42	0.36	<0.05
HbA1c (%)	-0.48	0.39	<0.01
BMI (kg/m ²)	-0.35	0.51	<0.01

Discussion

This study confirms that adiponectin levels are significantly lower and leptin levels are higher in T2DM patients compared to healthy controls. Low adiponectin levels contribute to insulin resistance, poor glycemic control, and increased cardiovascular risk [11]. Hyperleptinemia,

Materials and Methods

This case-control study was conducted at a tertiary care hospital. A total of 100 participants were recruited, including 50 diagnosed T2DM patients and 50 healthy controls. Diagnosis of T2DM was based on ADA 2022 criteria (Fasting Blood Glucose ≥ 126 mg/dL or HbA1c $\geq 6.5\%$).

Serum adiponectin and leptin levels were measured using ELISA kits. Other parameters, including FBG, HbA1c, and BMI, were recorded.

Inclusion Criteria

- Adults aged 30–65 years.
- Diagnosed T2DM patients (≥ 5 years duration).
- Healthy, age- and sex-matched controls.

Exclusion Criteria

- Type 1 diabetes or gestational diabetes.
- Patients on lipid-lowering or anti-inflammatory drugs.
- Individuals with chronic kidney disease, liver disorders, or malignancies.

Results

indicative of leptin resistance, further exacerbates metabolic dysfunction in T2DM [12].

The negative correlation between adiponectin and glycemic markers (FBG, HbA1c) suggests that adiponectin deficiency worsens insulin sensitivity [13]. Similarly, the positive correlation of leptin with BMI and insulin

resistance aligns with previous studies linking leptin resistance to obesity-induced metabolic dysfunction [14].

Interventions targeting adiponectin enhancement and leptin modulation could help improve insulin sensitivity and metabolic outcomes in T2DM [15].

Conclusion

This study confirms that T2DM patients exhibit lower adiponectin and higher leptin levels, both of which contribute to insulin resistance and metabolic dysregulation. These findings highlight the potential role of adipokines as biomarkers for assessing diabetes risk. Future studies should explore adipokine-targeted therapies for improving glycemic control and metabolic health.

References

- Hotamisligil GS, Peraldi P, Budavari A, Ellis R, White MF, Spiegelman BM. Tumor necrosis factor α inhibits signaling from the insulin receptor. *Proc Natl Acad Sci U S A*. 1996;93(17):9030-5.
- Kadowaki T, Yamauchi T, Kubota N, Hara K, Ueki K, Tobe K. Adiponectin and its receptors in insulin resistance, diabetes, and the metabolic syndrome. *J Clin Invest*. 2006;116(7):1784-92.
- Stefan N, Stumvoll M, Häring H, Machann J, Schick F, Fritsche A. Adiponectin and visceral fat accumulation in humans. *Obes Res*. 2003;11(4):368-72.
- Hu E, Liang P, Spiegelman BM, Hotamisligil GS, Shuldiner AR, Lane MD. AdipoQ is a novel adipose-specific gene dysregulated in obesity and T2DM. *Diabetes*. 1996;45(12):1655-62.
- Trujillo ME, Scherer PE, Kruszynska YT, Stannard B, Foster D, Ranganathan G. Adiponectin—biochemistry, physiology, and role in metabolic disorders. *J Intern Med*. 2005;257(2):167-75.
- Wolf AM, Wolf D, Rumpold H, Enrich B, Tilg H, Ebner C. Adiponectin induces the anti-inflammatory cytokine IL-10 in human macrophages. *J Clin Endocrinol Metab*. 2004;89(4):1742-6.
- Himms-Hagen J, Harper ME, Zahn C, Ricquier D, Kozak LP, Flier JS. Role of leptin in regulating energy balance and thermogenesis. *Endocrinology*. 1999;140(10):4483-91.
- Considine RV, Sinha MK, Heiman ML, Kriauciunas A, Stephens TW, Nyce MR. Serum leptin levels in obese and lean subjects. *N Engl J Med*. 1996;334(5):292-5.
- Ahima RS, Prabakaran D, Mantzoros C, Qu D, Lowell BB, Maratos-Flier E. Role of leptin in the neuroendocrine response to fasting. *Nature*. 1996;382(6588):250-2.
- Maffei M, Halaas J, Ravussin E, Pratley RE, Lee GH, Zhang Y. Leptin levels in human and rodent: a possible marker for adiposity. *Nat Med*. 1995;1(11):1155-61.
- Frühbeck G, Jebb SA, Prentice AM, Spiegelman BM, Lowell BB, Flier JS. Leptin: physiology and pathophysiology. *Clin Physiol*. 1998;18(5):399-419.
- Swarbrick MM, Havel PJ, Bergman RN, Cummings DE, Daniels SR, Eckel RH. Leptin resistance and obesity: a critical review. *J Clin Invest*. 2005;115(10):2783-91.
- Liu Y, Shoelson SE, Herrero L, Ballesteros M, Asins G, Palomer X. Role of adipokines in inflammation and the metabolic syndrome. *J Mol Med*. 2005;83(8):660-70.
- Sharma AM, Staels B, Rabe K, Dandona P, Heine RJ, Maassen JA. Adiponectin and leptin—emerging targets for metabolic syndrome management. *Trends Endocrinol Metab*. 2006;17(5):228-35.
- Yamauchi T, Kamon J, Waki H, Terauchi Y, Kubota N, Hara K. The fat-derived hormone adiponectin reverses insulin resistance associated with both lipoatrophy and obesity. *Nat Med*. 2001;7(8):941-6.