

RISK FACTORS OF DIABETIC FOOT ULCER AT A TERTIARY CARE HOSPITAL AMONG DIABETIC PATIENTS

Dr. Sanvar Mal Kantva¹, Dr. Mahendra Kumar²

¹ Senior Resident, Department of General Surgery, S M S Medical College & Attached Group of Hospital, Jaipur

² Assistant Professor, Department of General Surgery, Government Medical College Churu

Article Info: Received 20 July 2019; Accepted 18 August. 2019

DOI: <https://doi.org/10.32553/ijmbs.v3i8.480>

Address for Correspondence: Dr. Mahendra Kumar

Conflict of interest: No conflict of interest.

Abstract

Introduction: Diabetes is a metabolic disorder which affects not only carbohydrate but also protein and fat metabolism. Diabetes is also associated with acute as well chronic complications. Almost most of the organs, tissues and systems are affected due to long standing diabetes mellitus. Thus, adult and elderly with long standing diabetes patients often present to the clinics with multiple complications. Among all these well-known complications, diabetic foot ulcer is the most common. It affects as many as 15% of patients with diabetes mellitus during their lifetime

Material and Method: The study was a hospital based cross sectional study. Present study was carried out at outpatient department of General Surgery. After their verbal informed consent after explaining them the nature of the study. Patient confidentiality was maintained. The patients diagnosed with diabetic foot ulcer were given appropriate treatment, follow up and health education. The study patients were not subjected for any kind of invasive procedure for the present study purpose.

Results: 64.0% patients were having peripheral neuropathy, 73% were having >10 years of diabetes, 49% were having peripheral arterial disease, 34% patients were having diabetic Charcot joint & 26% were having uncontrolled hyperglycemia.

Conclusion: Diabetes Mellitus is a lifelong disease and diabetic foot complications can be life threatening, physically incapacitating, costly to treat and result in extensive morbidity.

Keywords: Diabetes, foot ulcers, neuropathy.

Introduction:

There are an estimated 42 million cases of diabetes mellitus in India. Thus, Indian ranks number in the top ten countries with high diabetes burden.^{1,2}

Diabetes is a metabolic disorder which affects not only carbohydrate but also protein and fat metabolism. Diabetes is also associated with acute as well chronic complications. Almost most of the organs, tissues and systems are affected due to long standing diabetes mellitus. Thus, adult and elderly with long standing diabetes patients often present to the clinics with multiple complications. Among all these well-known complications, diabetic foot ulcer is the most common. It affects as many as 15% of patients with diabetes mellitus during their lifetime.² Diabetic foot ulcer puts enormous social impact on the patient. The patient may get isolated socially, he may lose his prior social role, stigma and discrimination are common.³

Like other complications of diabetes mellitus, diabetic foot ulcer is also easily preventable. Proper care by diabetic patient has shown to reduce the incidence of diabetic foot ulcer as well as consequent foot amputation by about 80% which is a great advantage for the patient as well as society. Simple measures like control of blood sugar, appropriate diet, staying away from addictions like smoking, tobacco in any form, proper foot care, regular checkup, good hemoglobin levels, controlled blood pressure levels, controlled lipid levels etc.^{4,5}

To prevent the complications like diabetic foot, it is very essential that the patient should go for regular checkup and proper and adequate treatment.⁶

Hence present was undertaken to identify the risk factors and prevalence of diabetic foot ulcer in our present settings, so that patients can be made aware and can be managed adequately.

Materials & Method

The study was a hospital based cross sectional study. Present study was carried out at outpatient

department of General Surgery. After their verbal informed consent after explaining them the nature of the study. Patient confidentiality was maintained. The patients diagnosed with diabetic foot ulcer were given appropriate treatment, follow up and health education. The study patients were not subjected for any kind of invasive procedure for the present study purpose.

Sample size: 100 patients reporting to the Surgery dept. within study duration and eligible as per inclusion criteria

Inclusion Criteria: All patients of Diabetic foot who gave informed verbal consent

Exclusion Criteria:

Diabetic foot associated with venous ulcers and lymphedema.

Data thus collected were entered into excel and were then analyzed with help of EPI-Info software through tables, diagrams and appropriate statistical test wherever required.

RESULTS

Table 1: Distribution of cases according to socio-demographic variable

Variable	No of cases	Percentage
Male	57	57.00
Female	43	43.00
Mean age ± S D	59.23± 9.24 Yrs	
Hindu	79	79.00
Muslim	21	21.00

Maximum patients were hindu male and mean age of patients was 59.23± 9.24 Yrs.

Table 2: Distribution of cases according to risk factors responsible for ulceration:

RISK FACTOR	No.	(%)
Peripheral neuropathy	64	64.0
Foot Deformity	33	33.0
Diabetic Charcot Joint	34	34.0
Trauma	14	14.0
Improperly fitted shoes	13	13.0
Peripheral Arterial Disease	49	49.0
Callus	19	19.0
H/O Prior Ulceration	41	41.0
Higher Plantar Foot Pressure	21	21.0
Limited joint mobility	27	27.0
Uncontrolled hyperglycemia	26	26.0
Diabetes duration (>10 yrs)	73	73.0
Older Age	13	13.0
Poor Foot Hygiene	24	24.0

Above table shows that 64.0% patients were having peripheral neuropathy, 73% were having >10 years of diabetes, 49% were having peripheral arterial disease, 34% patients were having diabetic Charcot joint & 26% were having uncontrolled hyperglycemia.

DISCUSSION

In the present study, mean age was 59.23± 9.24 Yrs. Wheel Lock ⁷ did a study which revealed that the youngest age with diabetic foot was 32 years and the oldest age was 89 years. Mayfield et al ⁸ did a study on sex wise distribution of diabetic foot which included 32 males and 29 females.

In our study 64.0% patients were having peripheral neuropathy, 73% were having >10 years of diabetes, 49% were having peripheral arterial disease, 34% patients were having diabetic Charcot joint & 26% were having uncontrolled hyperglycemia.

Foot ulcers among diabetics are a major cause of mortality and morbidity and also a huge financial burden on health care services as inadequate and improper treatment could result in development of gangrene leading to limb amputations and sometimes even death. The significance lies in the fact that identification of risk factors is important to screen high-risk patients to prevent development of foot ulcers and its associated morbidity.⁹ The greatest risk factors for developing foot ulceration are neuropathy, arterial disease, foot deformity and peripheral neuropathy accounting for 80% of diabetic persons with foot ulcers. The contributing factors are poor glycemic control and microvascular disease.¹⁰

CONCLUSION

Diabetes Mellitus is a lifelong disease and diabetic foot complications can be life threatening, physically incapacitating, costly to treat and result in extensive morbidity.

REFERENCES

1. Ramachandran A, Ma RCW, Snehalatha C. Diabetes in Asia. Lancet. 2010 Jan 30;375(9712):408-18.
2. Shankhdhar K, Shankhdhar LK, Shankhdhar U, Shankhdhar S. Diabetic foot problems in India: an overview and potential simple approaches in a developing country. Curr Diab Rep. 2008 Dec 1;8(6):452-7.
3. Viswanathan V, Shobhana R, Snehalatha C, Seena R, Ramachandran A. Need for education on footcare in diabetic patients in India. J Assoc Physicians India. 1999 Nov;47(11):1083-5.

4. Shobhana R, Rama Rao P, Lavanya A, Viswanathan V, Ramachandra A. Cost burden to diabetic patients with foot complications-a study from southern India. JAPI. 2000;48(12):1147-50.
5. Harrington C, Zagari MJ, Corea J, Klitenic J. A cost analysis of diabetic lower-extremity ulcers. Diab Care. 2000 Sep 1;23(9):1333-8.
6. Croxson S. Diabetes in the elderly: problems of care and service provision. Diab Med. 2002 Jul;19:66- 72.
7. Wheel Lock, FC. Jr : Annals Of Surgery, 99:776:1969.
8. Mayfield, Jennifer A. A foot risk classification system to predict diabetic amputation in pima Indians. Diabetes care 1996, 704-09.
9. Thanh D, Aristidis V. The Influence of Gender as a Risk Factor in Diabetic Foot Ulceration. Available from: URL:<http://www.woundsresearch.com/article/8707>.
10. Vincent Lopez Rowe. Diabetic Ulcers. Available from: URL: <http://emedicine.medscape.com/article/460282-overview>