|| ISSN(online): 2589-8698 || ISSN(print): 2589-868X ||

International Journal of Medical and Biomedical Studies

Available Online at www.ijmbs.info

Volume 3, Issue 1; January: 2019; Page No. 73-78

PubMed (National Library of Medicine ID: 101738825)



Original Research Article

A STUDY ON SKELETALLY IMMATURE UNSTABLE PELVIC FRACTURES IN CHILDREN

Dr. Rakesh Kumar Gupta¹, Dr. Mohkam Singh²

¹Assistant Professor Dept. of Orthopaedics K.M. Medical College and Hospital, Mathura (UP).

²Assistant Professor Dept. of Orthopaedics K.M. Medical College and Hospital, Mathura (UP).

Article Info: Received 28 October 2018; Accepted 08 December. 2018

Cite this article as: Gupta, Dr. R. K., & Singh, Dr. M. (2019). A STUDY ON SKELETALLY IMMATURE UNSTABLE PELVIC FRACTURES IN CHILDREN. International Journal of Medical and Biomedical Studies, 3(1).

DOI: https://doi.org/10.32553/ijmbs.v3i1.602

Address for Correspondence: Dr. Mohkam Singh, Assistant Professor Dept. of Orthopaedics, K.M. Medical College and Hospital, Mathura (UP).

Conflict of interest: No conflict of interest.

Abstract

Introduction: Out of all factures the pelvic facture showed relatively less which account for 3%. In children Pelvic fractures are relatively rare as less than 0.2% of all paediatric fractures and 1-5% of admissions to tertiary children's health centre. Pelvis is a structure like a basin shaped bony which supports the spinal column and protects the abdominal organs. It may be facture due to high energy forces such as fall from a height or motor vehicle crash. Incident of pelvic fracture is increasing and these are associated with a high mortality rate of 25%. There was relatively low rate of occurrence of facture of pelvic injuries that associated with high levels of morbidity and mortality. According to the studied of the post-mortem examined only in children 66 deaths caused by trauma that showing pelvic fracture and severe bleeding to be the cause of death in 42% of the cases. In the case of adult in cases of unstable pelvic fractures retroperitoneal bleeding is the worst complication. When involving emergency situation cases with hemodynamic instability immediate treatment must be conducted using an external fixation, though it is only for temporarily to prioritise controlling the bleeding and saving the patient's life.

Aim: The main objective of this study is to study the surgical treatment for unstable pelvic fractures in children and stabilisation on basic primary care provided at an emergency service.

Material and methods: In this study total 10 children patients were included who had suffered unstable fractures of the pelvic ring. From all the patients detail clinical history were taken with analysis of the pre and postoperative radiographs by presence of the triradiate cartilage of the iliac. This study was based on radiographic quantification and the outcome of the residual pelvic and after the surgery treatment using the method of Keshishyan et alⁱ for comparison of pre and post operative findings.

Result: In this study unstable fractures of the pelvic ring evaluated where female were 7 and male were 3 with the ration 7: 3. The ranges were 2 to 13 years with the mean age 7.2 years old. For all the patients operation was done on an average of 12.3 days after the facture. The maximum cases the initial trauma were due to the run over followed by motorcycle accident and fall from height respectively. The injuries of the pelvis joint were divided inti anterior and posterior. Symphysis disjucction of the pubic facture were present in maximum with the facture of two rami and facture of four rami respectively. In five of the cases Sacroiliac dislocation correspond to posterior lesion were

seen. Unilateral fractures of the posterior ilium were also seen whereas unilateral anterior opening of the sacroiliac joint was also present.

Conclusion: In children facture of pelvic ring is rare and an indication for surgical treatment is unusual. Their concerns relate to the complications encountered that in leg length and residual pain in the sacroiliac joint. Hence for the justification, this study finding provides justifies the option of surgical treatment for reduction and correction of pelvic deformities, of the pelvic ring at an early stage, at the time of the injury.

Keywords: children, unstable pelvic facture, pelvic ring, Pelvic asymmetry

Introduction:

Out of all factures the pelvic facture showed relatively less which account for 3%ⁱⁱ. In children Pelvic fractures are relatively rare as less than 0.2% of all paediatric fractures and 1-5% of admissions to tertiary children's health centreⁱⁱⁱ, iv. These fractures are generally caused by highenergy trauma, high-impact accidents and are often associated with injuries to other organic lesions like abdominal viscera, genitourinary system, neurovascular, musculoskeletal structures and central nervous system^v.

Pelvis is a structure like a basin shaped bony which supports the spinal column and protects the abdominal organs. It may be facture due to high energy forces such as fall from a height or motor vehicle crash. Incident of pelvic fracture is increasing and these are associated with a high mortality rate of 25%^{vi}. There was relatively low rate of occurrence of facture of pelvic injuries that associated with high levels of morbidity and mortality.

According to the studied of the post-mortem examined only in children 66 deaths caused by trauma that showing pelvic fracture and severe bleeding to be the cause of death in 42% of the cases. In the case of adult in cases of unstable pelvic fractures retroperitoneal bleeding is the worst complication vii viii & ix. One of the studied showed that in 95 children suffering from pelvic fractures, locomotion capacity and dependence was assessed which showed that 80% of patients had unstable fractures and 52% of patients with stable fractures were dependent on help for locomotion^x. When involving emergency situation cases with hemodynamic instability immediate treatment must be conducted using

an external fixation, though it is only for temporarily to prioritise controlling the bleeding and saving the patient's life^{xi}. In cases with dislocation of the sacroiliac joint an anterior external fixator alone will not be sufficient to reduce and stabilise the posterior ring^{xii}.

The main objective of this study is to study the surgical treatment for unstable pelvic fractures in children and stabilisation on basic primary care provided at an emergency service.

MATERIAL AND METHODS:

This study was retrospective study carried out in the Dept. of orthopedics, K.M. Medical College and Hospital, Mathura (UP). India. In this study total 10 children patients were included who had suffered unstable fractures of the pelvic ring and had undergone surgical treatment in this institution during the period of 1 year. From all the patients detail clinical history were taken with analysis of the pre and postoperative radiographs by presence of the triradiate cartilage of the iliac. This study was based on radiographic quantification and the outcome of the residual pelvic and after the surgery treatment using the method of Keshishyan et alxiii for comparison of pre and post operative findings.

RESULT:

In this study total 10 patients were included with unstable fractures of the pelvic ring evaluated where female were 7 and male were 3 with the ration 7: 3. The age ranges were 2 to 13 years with the mean age 7.2 years old. For all the patients operation was done on an average of 12.3 days after the facture. In this study the maximum cases the initial trauma was due to the

run over followed by motorcycle accident and fall from height respectively. The injuries of the pelvis joint were divided inti anterior and posterior. Symphysis disjucction of the pubic facture were present in maximum with the facture of two rami and facture of four rami respectively. In five of the cases Sacroiliac dislocation correspond to posterior lesion were seen. Unilateral fracture of the posterior ilium were also seen whereas unilateral anterior opening of the sacroiliac joint was also present.

The AO-OTA classification was used to evaluate the cases with the following distribution. 61 B1 (one case); 61 B2 (one case); 61 C3 (one cases), 61 C2 (one case) and 61 C1 (seven cases) as shown in the table no 1 below. For all the

patients primary health care were provided before reached to our medical college and hospital after clinical stabilization was achieved. Six patients were suffered from other associated injuries as one with a clavicle fracture was treated without surgery using external immobilisation. Two with a fracture of the proximal humerus and fractures of the bones of the lower leg is ilateral to the pelvic lesion, one with the upper limb was treated conservatively and the tibia was fixed with flexible intramedullary nails. One with olecranon who suffered a fall was fixed using a tension band and one with concomitant injury to the bladder, diagnosed using exploratory laparotomy during primary care.

Table 1: showing the patients data with clinical presentation

	Age		AO-OTA			cause of
S.N.	(yrs)	sex	classification	posterior lesion	anterior lesion	accident
1	2	F	61C1	Sacroiliac dislocation	Disjunction of symphysis	Run over
						Motorcycle
2	12	F	61C1	Sacroiliac dislocation	Fracture of two rami	accident
3	5	М	61B1	Anterior sacroiliacopening	Fracture of two rami	Run over
4	5	F	61B2	Sacroiliacdislocation	Disjunction of symphysis	Run over
5	9	М	61C1	Sacroiliacdislocation	Disjunction of symphysis	Run over
6	10	F	61C1	Sacroiliacdislocation	Disjunction of symphysis	Run over
7	13	F	61C2	Sacroiliacdislocation +anterior compression (bilateral)		Motorcycle accident
8	6	М	61C1	Sacroiliacdislocation	Fracture of two rami	Run over
9	3	F	61C3	Sacroiliacdislocation (bilateral)	Fracture of four rami	Run over
						Fall from
10	7	F	61C1	Sacroiliacdislocation	Disjunction of symphysis	height

In the radiographic evaluation, pelvic asymmetry before the surgical procedure ranged from 0.7 to 2.8 cm (mean of 1.46 cm), and fall the range to 0.2 to 0.9 cm (mean 0.4 cm) after the reduction as shown in table no 2 below. The follow-up period for the patients in the hospital was ranged from 210 to 1,896 days.

Table 2: showing the result from surgery treatment

		Preoperative pelvic	Postoperative pelvic	Preoperative deformity index	Postoperative deformity index
S. N.	Surgery performed	asymmetry(cm	asymmetry (cm)	(Keshishyan)	(Keshishyan)
	Symphysis plate +				
1	sacroiliac screw	1.7	0.2	0.3	0.02
	External fixator +				
2	sacroiliac screw	0.7	0.3	0.05	0.02
	External fixator +				
3	sacroiliac screw	1.8	0.3	0.17	0.04

	External fixator +				
4	sacroiliac screw	1.1	0.5	0.2	0.04
	External fixator +				
5	sacroiliac screw	1.2	0.3	0.1	0.03
	External fixator +				
6	sacroiliac screw	1.6	0.5	0.1	0.05
	External fixator +				
7	sacroiliac screw	2.8	0.9	0.15	0.08
	External fixator +				
8	sacroiliac screw	0.9	0.2	0.12	0.05
	External fixator +				
9	sacroiliac screw	1.5	0.5	0.07	0.03
	External fixator +				
10	sacroiliac screw	1.2	0.2	0.08	0.03




Figure 1: Keshishyan et al. [1] method of measuring pelvic obliquity, where the pelvic asymmetry is the difference in length between the diagonals (X – Y cm). The degree of deformity corresponds to the difference in the diagonals (X – Y cm) divided by the sum of the diagonals (X + Y cm).



Figure 2: female, 12 years old, hit by a car, mechanism of trauma was lateral compression;

anterior injury was a fracture of four rami and posterior was a bilateral sacroiliac dislocation.



Figure 3: postoperative X-ray, AP view showing the final result.




Figure 4: postoperative X-ray, AP view showing supra-acetabular external fixation and cannulated screws in both sacroiliac joints.

DISCUSSION:

In children unstable pelvic fractures are relatively rare which are caused by high-impact traumas. In comparision to adults; in children Pelvic injuries are accompanied by various organ injuries that associated with high morbidity and are mortality^{xiv}, xv, xvi xvii & xviii. In this study run over on public roads was main cause. The profile of the children was reported to be literature excepted for the distribution according to sex. However, most frequently lesion occur in male gender but whereas in this study maximum cases were female almost 70 %. In this study run over on public roads was the most frequent cause of the fracture that was agree with the studied done by Signorino et al^{xix} who reported as commonest causes were being run over and traffic accidents.

There are many studied which have analyzed the long-term evolution of unstable and stable lesions of the pelvic ring that were treated conservatively^{xx}, xxi & xxii. Studied of Schwarz et al^{xxiii}. showed that 47% rate of poor results due to pelvic asymmetry; which underline the importance of achieving anatomical reduction and conclude that unstable pelvic fractures. Several studies presented long-term progression of unstable fractures of the pelvic ring that were non-surgically and treated evolved unsatisfactory results, with residual pain in the posterior region of pelvis which is due to residual pelvic asymmetry. None of the cases treated surgically whereas in this studied showed evolved with residual pain, step on abnormality on physical examination xxiv, xxv & xxvi.

In this study showed the limitation with the sample size this is because of the rare fracture that affects around one in 100,000 children per year and we believe sample is relevant, since no large published series on surgical treatment for this lesion exist. The result of this study helps to encourage continue to indicate surgical treatment for cases in which there is pelvic asymmetry greater than 0.5 cm when there is present of associated dislocation of the sacroiliac joint, that would be unlikely to evolve satisfactorily without surgical measures.

The combination with Posterior fixation with a sacroiliac cannulated screw in the sacroiliac joint was based on the facts that this technique is noninvasive, that provides adequate mechanical stability to the posterior portion of the pelvis has been shown to be a safe and reproducible method in children. Therefore the application of an anterior external fixator does not enable the control of the posterior elements of the pelvic ring xxvii.

CONCLUSION:

In children facture of pelvic ring is rare and an indication for surgical treatment is unusual. There are many research which still have a controversy for treatment, which has been recommended for treating these fractures for many years. Their concerns relate to the complications encountered that in leg length and residual pain in the sacroiliac joint. Hence for the justification, this study finding provides justifies the option of surgical treatment for reduction and correction of pelvic deformities, of the pelvic ring at an early stage, at the time of the injury.

REFERENCES:

- 1. Keshishyan RA, Rozinov VM, Malakhov AO, Kuznetsov LE, Strunin EG, Cho-govadze GA, et al. Pelvic polyfractures in children. Radiographic diagnosis and treatment. Clin Orthop Relat Res. 1995;(320):28-33
- **2.** Bailey & Love: Short practice of surgery. 24th Edition international Edition Arnold publishers London 2004:387-388.
- **3.** Smith WR, Oakley M, Morgan SJ. Pediatric pelvic fractures. J Pediatr Orthop 2004; 24:130–5.
- **4.** Chia JP, Holland AJ, Little D, Cass DT. Pelvic fractures and associated injuries in children. J Trauma 2004;56:83–8.
- 5. Smith W, Shurnas P, Morgan S, Agudelo J, Luszko G, Knox EC, et al. Clinical outcomes of unstable pelvic fractures in skeletally immature patients. J Bone Joint Surg Am. 2005;87(11):2423-31

- **6.** Rittmeister M, Lindsey RW, Kohl HW: Pelvic fracture among polytrauma decedents. Trauma based mortality with pelvic fracture a case series of 74 patients. 3rd. The ben Taub general hospital, Baylor College of medicine, Houston, U S A. Arch Ortho trauma surg. 2001; 121(1):43-9
- Signorino PR, Densmore J, Werner M, Winthrop A, Stylianos S, Guice KS, et al. Pediatric pelvic injury: functional outcome at 6-month follow-up. J Pediatr Surg 2005;40: 107–13
- **8.** Keshishyan RA, Rozinov VM, Malakhov AO, Kuznetsov LE, Strunin EG, Chogovadze GA, et al. Pelvic polyfractures in children. radiographic diagnosis and treatment. Clin Orthop Rel Res 1995;(320):28–33.
- Upperman JS, Gardner M, Gaines B, Schall L, Ford HR. Early functional outcome in children with pelvic fractures. J Pediatr Surg 2000;35:1002–5
- **10.** Blasier DR, McAtee J, White R, Mitchell DT. Disruption of the pelvic ring in pediatric patients. Clin Orthop Relat Res. 2000; (376):87-95.
- **11.** Cothren CC, Moore EE, Smith WR, Morgan SJ. Preperitoneal pelvic packing in the child with an unstable pelvis: a novel approach. J Pediatr Surg 2006;41:e17–9.
- **12.** Quick TJ, Eastwood DM. Pediatric fractures and dislocations of the hip and pelvis. Clin Orthop Rel Res 2005;432:87–96.
- **13.** Keshishyan RA, Rozinov VM, Malakhov AO, Kuznetsov LE, Strunin EG, Cho-govadze GA, et al. Pelvic polyfractures in children. Radiographic diagnosis and treatment. Clin Orthop Relat Res. 1995;(320):28-33
- **14.** Blasier RD, McAtee J, White R, Mitchell DT. Disruption of the pelvic ring in pediatric patients. Clin Orthop Rel Res 2000;376:87–95.
- **15.** Banerjee S, Barry MJ, Paterson JM. Paediatric pelvic fractures: 10 years' experience in a trauma centre. Injury 2009;40:410–3.
- **16.** Grisoni N, Connor S, Marsh E, Thompson GH, Cooperman DR, Blakemore LC. Pelvic

- fractures in a pediatric level I trauma center. J Orthop Trauma 2002;16:458–63.
- **17.** Heeg M, Klasen HJ. Long-term outcome of sacroiliac disruption in children. J Pediatr Orthop 1997;17:337–41.
- **18.** McDonald GA. Pelvic disruptions in children. Clin Orthop Rel Res 1980;151:130–4.
- **19.** Signorino PR, Densmore J, Werner M, Winthrop A, Stylianos S, Guice KS, et al. Pediatric pelvic injury: functional outcome at 6-month follow-up. J Pediatr Surg. 2005;40(1):107-13
- **20.** Heeg M, Klasen HJ. Long-term outcome of sacroiliac disruption in children. J Pediatr Orthop. 1997;17(3):337-41.
- **21.** McDonald GA. Pelvic disruptions in children. Clin Orthop Relat Res. 1980;(151):130-4
- **22.** Bryan WJ, Tullos HS. Pediatric pelvic fractures: review of 52 patients. J Trauma. 1979;19(11):799-805
- **23.** Schwarz N, Posch E, Mayr J, Fischmeister FM, Schwarz AF, Öhner T. Long-term results of unstable pelvic ring fractures in children. Injury 1998;29:431–3.
- **24.** Smith W, Shurnas P, Morgan S, Agudelo J, Luszko G, Knox EC, et al. Clinical outcomes of unstable pelvic fractures in skeletally immature patients. J Bone Joint Surg Am. 2005;87(11):2423-31.
- **25.** McDonald GA. Pelvic disruptions in children. Clin Orthop Relat Res. 1980;(151):130-4.
- **26.** Quick TJ, Eastwood DM. Pediatric fractures and dislocations of the hip and pelvis. Clin Orthop Rel Res. 2005;(432):87-96.
- **27.** Matta JM, Anderson LM, Epstein HC, Hendrick P. Fractures of the acetabulum. A retrospective analysis. Clin Orthop Relat Res. 1986;(205):230-40.